

A unified BFKL/DGLAP description of Deep Inelastic Scattering

J. Kwieciński^{*,†}, A.D. Martin[†] and A.M. Stasto^{*,†}

^{*} *H. Niewodniczański Institute of Nuclear Physics, Department of Theoretical Physics,
ul. Radzikowskiego 152, 31-342 Krakow, Poland.*

[†] *Department of Physics, University of Durham, Durham, DH1 3LE, UK.*

(presented by R.G. Roberts)

Abstract. We introduce a coupled pair of evolution equations for the unintegrated gluon distribution and the sea quark distribution which incorporate both the resummed leading $\ln(1/x)$ BFKL contributions and the resummed leading $\ln(Q^2)$ DGLAP contributions. We solve these unified equations in the perturbative QCD domain. With only two physically motivated parameters we obtain an excellent description of the HERA F_2 data.

1. INTRODUCTION

One of the striking features of the measurements of deep inelastic scattering at HERA is the strong rise of the proton structure function F_2 as x decreases from 10^{-2} to below 10^{-4} [1]. At first sight it appeared that the rise was due to the (BFKL) resummation of leading $\ln(1/x)$ contributions. In this approach the basic dynamical quantity at small x is the gluon distribution $f(x, k_T^2)$ *unintegrated* over its transverse momentum k_T . Observables are computed in terms of f via the k_T factorization theorem. For example

$$F_2 = F_2^{\gamma g} \otimes f \quad \text{where} \quad F_2^{\gamma g} = \sum_q e_q^2 B_q \quad (1)$$

and where $\otimes$ denotes a convolution in transverse (k_T), as well as longitudinal, momentum, see Fig. 1(a). $F_2^{\gamma g}$ is the off-shell gluon structure function which at lowest order is given by the “quark box and crossed-box” contributions, $\gamma g \rightarrow q\bar{q}$. On the other hand the conventional DGLAP approach is based on collinear factorization

$$\frac{\partial F_2}{x \partial \ln(Q^2/\Lambda^2)} = \sum_i e_i^2 P_{qg} \otimes g + \sum_i e_i^2 P_{qq} \otimes (q_i + \bar{q}_i) \quad (2)$$




FIGURE 1. (a) The diagrammatic representation of the k_T factorization formula $F_2 = F_2^{\gamma g} \otimes f$. At lowest order $F_2^{\gamma g}$ is given by the quark box shown (together with the crossed box) summed over all types of quark. (b) The different regions of integration used to evaluate the sea quark distribution. In regions II and III the sea is driven by the gluon, $S_q = B_q \otimes f$.

where $\overline{\otimes}$ is simply a convolution over longitudinal momentum. DGLAP evolution effectively sums up the leading $\ln Q^2$ contributions, and is able to describe the F_2 data at the smallest x values observed (even for $Q^2 \sim 1\text{GeV}^2$) with an appropriate choice of input parton distributions. These non-perturbative input shapes in x mean that there is more freedom in the pure DGLAP description than in the BFKL approach. In the latter approach, the shape in x emerges, in principle, from the perturbative $\ln(1/x)$ resummation. In practice it is not so clear cut since non-perturbative, and also subleading $\ln(1/x)$, effects can modify the BFKL prediction. Here we should attempt to find a unified description of F_2 which incorporates both (DGLAP and BFKL) of these perturbative effects.

2. UNIFIED BFKL/DGLAP FORMALISM

We argue that the unintegrated gluon distribution $f(x, k_T^2)$ and the k_T factorization theorem provides the natural framework for describing observables at small x . To determine f we arrange the BFKL equation so that we only need to solve it in the perturbative domain $k_T^2 > k_0^2$ [2]. We also include the residual DGLAP contributions. To be precise we have

$$f(x, k_T^2) = \frac{\alpha_S}{2\pi} \int_x^1 dz P_{gg}(z) \frac{x}{z} g\left(\frac{x}{z}, k_0^2\right) +$$

$$+ \frac{3\alpha_S}{\pi} k_T^2 \int_x^1 \frac{dz}{z} \int_{k_0^2}^{\frac{dk_T'^2}{k_T^2}} \left[\frac{f\left(\frac{x}{z}, k_T'^2\right) \theta\left(\frac{k_T^2}{z} - k_T'^2\right) - f\left(\frac{x}{z}, k_T^2\right)}{|k_T'^2 - k_T^2|} + \frac{f\left(\frac{x}{z}, k_T^2\right)}{(4k_T'^4 + k_T^4)^{\frac{1}{2}}} \right]$$

$$+ \frac{3\alpha_S}{\pi} \int_x^1 dz \left(\frac{P_{gg}(z)}{6} - \frac{1}{z} \right) \int_{k_0^2}^{k_T^2} \frac{dk_T'^2}{k_T'^2} f\left(\frac{x}{z}, k_T'^2\right) + \frac{\alpha_S}{2\pi} \int_x^1 dz P_{gq} \Sigma\left(\frac{x}{z}, k_T^2\right) \quad (3)$$

where $-1/z$ is taken from DGLAP because it is already included in BFKL. The input term comes from two sources: the $k_T^2 < k_0^2$ parts of BFKL and DGLAP terms. We specify the input in terms of a simple two parameter form $g(x, k_0^2) = N(1-x)^\beta$. In addition to restricting the solution of the BFKL equation to the perturbative region $k_T^2 > k_0^2$ and to including the DGLAP terms, we have also introduced a θ function which imposes the constraint $k_T'^2 < k_T^2/z$ on the real gluon emissions. The origin of this constraint¹ is the requirement that the virtuality of the exchanged gluon is dominated by its transverse momentum $|k'|^2 \simeq k_T'^2$. We take a running coupling $\alpha_S(k_T^2)$, which is supported by the results of the next-to-leading order $\ln(1/x)$ analyses of Fadin, Lipatov, Camici and Ciafaloni. The final term in (3) depends on the quark singlet momentum distribution Σ . At small x the sea quark components S_q of Σ dominate. They are driven by the gluon via the $g \rightarrow q\bar{q}$ transitions, that is $S_q = B_q \otimes f$ where at lowest order B_q is the box (and crossed box) contribution indicated in Fig. 1(a). Besides the z and k_T^2 integrations symbolically denoted by $\otimes$ the box contribution implicitly includes an integration over the transverse momentum κ_T of the exchanged quark. The evolution equation for Σ may be written in the form

$$\Sigma = S^{(0)} + \sum_q B_q(k_T^2 = 0) \otimes z g(z, k_0^2) + \sum_q B_q \otimes f + P_{qq} \otimes S_q + V \quad (4)$$

where the first three terms on the right hand side are the “ $B_q \otimes f$ ” contributions coming from the regions I, II, III of the k_T^2 and κ_T^2 integrations that are shown in Fig. 1(b). First, in the non-perturbative domain, region I, the u, d, s sea quark contribution is parametrized in the form $S^{(0)} = C_P x^{-0.08} (1-x)^8$ consistent with soft pomeron and counting rule expectations, where C_P is independent of Q^2 . The constant C_P is fixed in terms of the two parameters, N and β , by the momentum sum rule. In region II we apply the strong k_T ordering approximation with $B_q \approx B_q(k_T^2 = 0)$ so that the k_T^2 integration can be carried out to give a contribution proportional to $g(x/z, k_0^2)$. Finally in region III we evaluate the full box contribution; this gives the main contribution and is responsible for the rise of F_2 with decreasing x . The last two terms in (4) give the sea $\rightarrow$ sea evolution contribution, and the valence contribution $V(x, Q^2)$ which is taken directly from a recent parton set. The charm quark

¹⁾ A more general treatment of the gluon ladder which incorporates both the BFKL equation and DGLAP evolution is given by the CCFM equation [3], which is based on angular ordering of the gluon emissions. The angular ordered and kinematic constraints lead to similar subleading $\ln(1/x)$ effects, but the kinematic constraint overrides the angular ordered constraint, except when $Q^2 < k_T^2$ in the large x domain [4].




FIGURE 2. The two-parameter description of some of the F_2 data at small x using $f(x, k_T^2)$ evaluated with (continuous curves) and without (dashed curves) the kinematic constraint $k_T'^2 < k_T^2/z$.

component of the sea is given totally by *perturbative* QCD, since for $k_T^2 < k_0^2$ the box $B(k_T^2 = 0)$ is finite as $\kappa_T^2 \rightarrow 0$ due to $m_c \neq 0$.

3. DESCRIPTION OF F_2 DATA AND DISCUSSION

We solve the coupled integral equations (3) and (4) for the gluon f and the quark singlet Σ in the perturbative domain, $k_T^2 > k_0^2$. The only input is the gluon $g(x, k_0^2)$. We take $k_0^2 = 1\text{GeV}^2$. We determine the values of the two input parameters by fitting to the available data [1] for F_2 with $x < 0.05$ and $Q^2 > 1.5\text{GeV}^2$. The continuous curves in Fig. 2 show the description of a sample of the data. Overall the fit is excellent; at least as good as that achieved in the recent global analyses. When we repeat the analysis with the kinematic constraint omitted we see that the description (given by the dashed curves in Fig. 2) is not so good and, moreover, the extrapolation of the gluon to $x \approx 0.4$ no longer describes the WA70 prompt photon data. How important are the $\ln(1/x)$ effects? First we replace the BFKL kernel in (3) by the standard DGLAP splitting function, P_{gg} . We find that the gluon is not changed by much in the HERA domain $x \gtrsim 10^{-4}$, compare the dashed and dotted curves in Fig. 3. This effect is well known. In the power series expansion of the gluon anomalous dimension in α_S/ω the coefficients of the 2nd, 3rd and 5th terms are zero, where ω is the moment variable. On the other hand, when we use pure DGLAP evolution for the quark singlet, as well as the gluon, the difference is pronounced; compare the dashed and dot-dashed




FIGURE 3. The continuous and dashed curves come from solving (3) and (4) with and without the kinematic constraint. The dotted and dot-dashed curves are obtained using DGLAP in the gluon sector and in both the gluon and quark sectors respectively. The same input is used for all curves.

curves in Fig. 3.

We would like to conclude that we have a theoretically well-grounded and consistent formalism which, with the minimum of non-perturbative input, is able to give a good *perturbative* description of the observed structure of F_2 . Moreover the BFKL/DGLAP components of F_2 are decided by dynamics. In this way we have made a determination of the *universal* gluon distribution $f(x, k_T^2)$ which can be used, via the k_T factorization theorem, to predict the behaviour of other observables at small x . The predictions for F_2 (charm) and F_L can be found in [2].

REFERENCES

1. H1 Collab., S. Aid et al., *Nucl. Phys.* **B420** (1996) 3; ZEUS Collab., M. Derrick et al., *Z. Phys.* **C69** (1996) 607; *Z. Phys.* **C72** (1996) 399. BCDMS collaboration: A.C. Benvenuti et al., *Phys. Lett.* **B223** (1989) 485; New Muon collaboration (NMC): M. Arneodo et al., *Nucl. Phys.* **B483** (1997) 3; E665 collaboration: M.R. Adams et al., *Phys. Rev.* **D54** (1996) 3006.
2. J. Kwiecinski, A.D. Martin and A.M. Stasto, Durham preprint DTP/97/18.
3. M. Ciafaloni, *Nucl. Phys.* **B296** (1988) 49; S. Catani, F. Fiorani and G. Marchesini, *Phys. Lett.* **B234** (1990) 339; *Nucl. Phys.* **B336** (1990) 18.
4. J. Kwiecinski, A.D. Martin and P.J. Sutton, *Z. Phys.* **C71** (1996) 585; B. Andersson, G. Gustafson, H. Kharraziha and J. Samuelson *Z. Phys.* **C71** (1996) 613.